CONTENTS

Special Issue: QSAR in Environmental Toxicology. Proceedings of the Fourth International Workshop, Veldhoven, Netherlands, 16–20 September 1990

Preface xiii
List of Participants xv

Introductory Papers
QSAR in environmental sciences and drug design
J. Hermens (Utrecht, Netherlands). 1
Design of site-specific protein drugs
E. Tomlinson (Cambridge, MA, USA) 9
Structure–activity relationships of chemical mutagens and carcinogens
C. Hansch (Claremont, CA, USA). 17

Section 1: QSAR Techniques and Molecular Descriptors
Lipophilicity indices of triazine herbicides
G.L. Biagi, M.C. Guerra, A.M. Barbaro, M. Recanatini (Bologna, Italy), P.A. Borea (Ferrara, Italy) and A. Sapone (Bologna, Italy) 33
A validation study for the estimation of aqueous solubility from n-octanol/water partition coefficients
R. Brüggemann and J. Altschuh (Neuherberg, FRG) 41
The QSAR prediction of melting point, a property of environmental relevance
J.C. Dearden (Liverpool, UK) 59
Retention of neutral and basic heteroaromatic hydrocarbons in RPLC systems and its use in predictive studies. I. Concentration of the organic modifier
P. de Voogt, J.W.M. Wegener, U.A.Th. Brinkman and H. Govers (Amsterdam, Netherlands) 69
A thermodynamic analysis of quantitative structure–activity relationships for hydrophobic organic chemicals
F.A.P.C. Gobas (Windsor, Ont., Canada). 89
Descriptors for isomer resolution of (bio-)distribution of chlorinated aromatic compounds
H.A.J. Govers, R. Luijk and E.H.G. Evers (Amsterdam, Netherlands) . 105
Calculating the hydrophobicity of chlorinated hydrocarbon solutes
A.J. Leo (Claremont, CA, USA) 121
Outliers: their origin and use in the classification of molecular mechanisms of toxicity
R.L. Lipnick (Washington, DC, USA) 131
Complex factors in hydrocarbon/water, soil/water and fish/water partitioning
P.S. Magee (Vallejo, CA, USA) 155
Evaluation of some hydrophobic aspects of polymethyl- and polychloro-derivatives of benzene and biphenyl
R.F. Rekker, G. de Vries and R.E. Koopmans (Amsterdam, Netherlands) 179

Chemical topology and ecotoxicology
A. Sabljić (Zagreb, Croatia, Yugoslavia) 197
Do Hammett constants model electronic properties in QSARs?
G. Schüürmann (Leverkusen, FRG) 221
Prediction of contradictions between technologically and environmentally relevant properties of PCBs
A. van Haelst and H. Govers (Amsterdam, Netherlands) 237
Vapor pressure estimation for organic compounds
S.H. Yalkowsky (Tucson, AZ, USA) and D.S. Mishra (Pittsburgh, PA, USA) 243

Section 2: Environmental Fate Processes
Predictive QSAR models for estimating biodegradation of aromatic compounds
P. Degner, M. Nendza and W. Klein (Schmallenberg-Grafschaft, FRG) 253
Estimating atmospheric degradation processes by SARs
M. Müller and W. Klein (Schmallenberg-Grafschaft, FRG) 261
QSARs and PARs for biodegradation of PCBs
J.R. Parsons, L.C.M. Commandeur, H.E. van Eyseren and H.A.J. Govers (Amsterdam, Netherlands) 275
QSARs for predicting biotic and abiotic reductive transformation rate constants of halogenated hydrocarbons in anoxic sediment systems
W.J.G.M. Peijnenburg, M.J. 't Hart, H.A. den Hollander, D. van de Meent, H.H. Verboom (Bilthoven, Netherlands) and N.L. Wolfe (Athens, GA, USA) 283
Application of QSAR to biodegradation of linear alkylbenzene sulphonate (LAS) isomers and homologues
D.W. Roberts (Wirral, UK) 301
Multivariate modelling of the rate constant of the gas-phase reaction of haloalkanes with the hydroxyl radical
M.L. Tosato (Roma, Italy), C. Chiorboli (Ferrara, Italy), L. Eriksson and J. Jonsson (Umeå, Sweden) 307
QSAR analyses of oxidation and reduction rates of environmental organic pollutants in model systems
P.G. Tratnyek, J. Hoigné, J. Zeyer and R.P. Schwarzenbach (Dübendorf, Switzerland) 327
Predictive QSPR models for estimating soil sorption coefficients: potential and limitations based on dominating processes
B. von Oepen, W. Kördel, W. Klein and G. Schüürmann (Schmallenberg, FRG) 343

Section 3: Biokinetics and Bioaccumulation
A kinetic description of the fate of chemicals in biosystems
S. Baláž (Bratislava, Czechoslovakia), M. Wiese and J.K. Seydel (Borstel, FRG) 357

Kinetics of bioaccumulation and clearance of isomeric hexachlorocyclohexanes
W. Butte, K. Fox and G.-P. Zauke (Oldenburg, FRG) 377
Toxicokinetics of aromatic amines in guppy, *Poecilia reticulata*
W. de Wolf, E. Yedema, W. Seinen and J.L.M. Hermens (Utrecht, Netherlands) 383
QSAR for organic chemical bioconcentration in *Daphnia*, algae, and mussels
H.J. Geyer, I. Scheunert, R. Brüggemann, C. Steinberg (Neuherberg, FRG), F. Korte (Freising-Weihenstephan, FRG) and A. Kettrup (Neuherberg, FRG) 387
First-order and pseudo-first-order elimination kinetics
G. Schüürmann (Leverkusen, FRG) 395
QSAR approach to target organ estimation
M. Tichý (Prague, Czechoslovakia) 407
Bioconcentration, metabolism and toxicity of substituted anilines in the zebrafish (*Brachydanio rerio*)
S. Zok, G. Görge, W. Kalsch and R. Nagel (Mainz, FRG) 411

Section 4: Toxicity and Mechanism of Action
Correlation of tumour-inhibiting activity with acute toxicity
S.J.A. Bradburne and J.C. Dearden (Liverpool, UK) 425
QSAR studies of comparative toxicity in aquatic organisms
M.T.D. Cronin, J.C. Dearden (Liverpool, UK) and A.J. Dobbs (Marlow, UK) 431
Qualitative and quantitative modelling of toxic effects of organophosphorous compounds to fish
J. de Bruijn and J. Hermens (Utrecht, Netherlands) 441
Influence of survival time on the lethal body burden of 2,3,4,5-tetrachloroaniline in the guppy, *Poecilia reticulata*
W. de Wolf, A. Opperhuizen, W. Seinen and J.L.M. Hermens (Utrecht, Netherlands) 457
QSARs for the sublethal responses of marine mussels (*Mytilus edulis*)
P. Donkin, J. Widdows, S.V. Evans and M.D. Brinsley (Plymouth, UK) 461
Inorganic concepts relevant to metal binding, activity, and toxicity in a biological system
J.D. Hoeschele (Ann Arbor, MI, USA), J.E. Turner and M.W. England (Oak Ridge, TN, USA) 477
QSARs in phytotoxicity
E.M. Hulzebos (Bilthoven, Netherlands), D.M.M. Adema (Delft, Netherlands), E.M. Dirven-van Breemen (Bilthoven, Netherlands), L. Henzen (Delft, Netherlands) and C.A.M. van Gestel (Bilthoven, Netherlands) 493
Regression and cluster analysis of the acute toxicity of 267 chemicals to six species of biota and the octanol/water partition coefficient
K.L.E. Kaiser and S.R. Esterby (Burlington, Ont., Canada) 499

Interpreting aquatic toxicity QSARs: the significance of toxicant body residues at the pharmacologic endpoint
L.S. McCarty (Mississauga, Ont., Canada), D. Mackay (Toronto, Ont., Canada), A.D. Smith, G.W. Ozburn (Thunder Bay, Ont., Canada) and D.G. Dixon (Waterloo, Ont., Canada) 515
Predictive QSAR models for estimating ecotoxic hazard of plant-protecting agents: target and non-target toxicity
M. Nendza, B. Dittrich, A. Wenzel and W. Klein (Schmallenberg-Grafschaft, FRG) 527
A QSAR study of the toxicity of amines to the fathead minnow
L.D. Newsome, D.E. Johnson, R.L. Lipnick (Washington, DC, USA), S.J. Broderius and C.L. Russom (Duluth, MN, USA). 537
The utility of computed superdelocalizability for predicting the LC_{50} values of epoxides to guppies
R. Purdy (St. Paul, MN, USA) 553
QSAR issues in aquatic toxicity of surfactants
D.W. Roberts (Wirral, UK) 557
QSARs for monosubstituted anilines eliciting the polar narcosis mechanism of action
T.W. Schultz, D.T. Lin and L.M. Arnold (Knoxville, TN, USA) 569
QSARs for selected aliphatic and aromatic amines
T.W. Schultz, T.S. Wilke, S.E. Bryant and L.M. Hosein (Knoxville, TN, USA) 581
Development of QSARs in terrestrial ecotoxicology: earthworm toxicity and soil sorption of chlorophenols, chlorobenzenes and dichloroaniline
C.A.M. van Gestel (Bilthoven, Netherlands), W.-c. Ma (Arnhem, Netherlands) and C.E. Smit (Bilthoven, Netherlands) 589
QSARs for toxicity of organophosphorous pesticides to *Daphnia* and honeybees
M. Vighi, M.M. Garlanda and D. Calamari (Milano, Italy) 605

Section 5: Application of QSAR in Practice

Data base of aqueous solubility for organic non-electrolytes
R.-M. Dannenfelser and S.H. Yalkowsky (Tucson, AZ, USA) 625
The use of data estimation methods by regulatory authorities
J.W. Hart (Copenhagen, Denmark) 629
Development of a predictive model for biodegradability based on BIODEG, the evaluated biodegradation data base
P.H. Howard (Syracuse, NY, USA), R.S. Boethling (Washington, DC, USA), W. Stiteler, W. Meylan and J. Beauman (Syracuse, NY, USA) . . 635
Performance of on-line chemical property estimation methods with TSCA premanufacture notice chemicals
D.G. Lynch, N.F. Tirado, R.S. Boethling (Washington, DC, USA), G.R. Huse (Springfield, VA, USA) and G.C. Thom (Washington, DC, USA) 643

Environmental hazard and risk assessment under the United States Toxic Substances Control Act
J.V. Nabholz (Washington, DC, USA) 649
ASTER: an integration of the AQUIRE data base and the QSAR system for use in ecological risk assessments
C.L. Russom, E.B. Anderson, B.E. Greenwood and A. Pilli (Duluth, MN, USA). 667
QSARs in the chemical industry: need, scope and current limitations
G. Schüürmann and M. Marsmann (Leverkusen, FRG). 671
QSARs in Netherlands water quality management policies
M.A. van der Gaag (Lelystad, Netherlands) 677
The application of QSARs, extrapolation and equilibrium partitioning in aquatic effects assessment for narcotic pollutants
C.J. van Leeuwen, P.T.J. van der Zandt (Leidschendam, Netherlands), T. Aldenberg (Bilthoven, Netherlands), H.J.M. Verhaar and J.L.M. Hermens (Utrecht, Netherlands). 681
Chemical selection by the Interagency Testing Committee: use of computerized substructure searching to identify chemical groups for health effects, chemical fate and ecological effects testing
J.D. Walker (Washington, DC, USA) 691

Author Index 701
Subject Index 703